

A DICHOTOMY PROPERTY FOR LOCALLY COMPACT GROUPS

MARITA FERRER, SALVADOR HERNÁNDEZ, AND LUIS TÁRREGA

Respectfully dedicated to Professor Wis Comfort

ABSTRACT. We extend to metrizable locally compact groups Rosenthal's theorem describing those Banach spaces containing no copy of ℓ_1 . For that purpose, we transfer to general locally compact groups the notion of interpolation (I_0) set, which was defined by Hartman and Ryll-Nardzewsky [25] for locally compact abelian groups. Thus we prove that for every sequence $\{g_n\}_{n<\omega}$ in a locally compact group G , then either $\{g_n\}_{n<\omega}$ has a weak Cauchy subsequence or contains a subsequence that is an I_0 set. This result is subsequently applied to obtain sufficient conditions for the existence of Sidon sets in a locally compact group G , an old question that remains open since 1974 (see [32] and [20]). Finally, we show that every locally compact group strongly respects compactness extending thereby a result by Comfort, Trigos-Arrieta, and Wu [13], who established this property for abelian locally compact groups.

1. INTRODUCTION

A well-known result of Rosenthal establishes that if $\{x_n\}_{n<\omega}$ is a bounded sequence in a Banach space X , then either $\{x_n\}_{n<\omega}$ has a weak Cauchy subsequence, or $\{x_n\}_{n<\omega}$ has a subsequence equivalent to the usual ℓ_1 -basis (and then X contains a copy of ℓ_1). In this paper, we look at this result for locally compact groups. More precisely, our main goal is to extend Rosenthal's dichotomy theorem on Banach spaces to locally compact groups and their weak topologies. First, we need some definitions and basic results.

Date: April 12, 2017.

Research Partially supported by the Spanish Ministerio de Economía y Competitividad, grant MTM2016-77143-P, and the Universitat Jaume I, grant P1171B2015-77. The second author also acknowledges partial support by Generalitat Valenciana, grant code: PROMETEO/2014/062.

2010 Mathematics Subject Classification. Primary 22D05; 43A46. Secondary 22D10; 22D35; 43A40; 54H11

Key Words and Phrases: Locally compact group; I_0 set; Sidon set; Interpolation set; Weak topology; Bohr compactification; Eberlein compactification.

Given a locally compact group (G, τ) , we denote by $Irr(G)$ the set of all continuous unitary irreducible representations σ defined on G . That is, continuous in the sense that each matrix coefficient function $g \mapsto \langle \sigma(g)u, v \rangle$ is a continuous map of G into the complex plane. Thus, fixed $\sigma \in Irr(G)$, if \mathcal{H}^σ denotes the Hilbert space associated to σ , we equip the unitary group $\mathbb{U}(\mathcal{H}^\sigma)$ with the weak (equivalently, strong) operator topology. For two elements π and σ of $Irr(G)$, we write $\pi \sim \sigma$ to denote the relation of unitary equivalence and we denote by \widehat{G} the *dual object* of G , which is defined as the set of equivalence classes in $(Irr(G)/\sim)$. We refer to [14, 4] for all undefined notions concerning the unitary representations of locally compact groups.

Adopting, the terminology introduced by Ernest in [16], set $\mathcal{H}_n \stackrel{\text{def}}{=} \mathbb{C}^n$ for $n = 1, 2, \dots$; and $\mathcal{H}_0 \stackrel{\text{def}}{=} l^2(\mathbb{Z})$. The symbol $Irr_n^C(G)$ will denote the set of irreducible unitary representations of G on \mathcal{H}_n , where it is assumed that every set $Irr_n^C(G)$ is equipped with the compact open topology. Finally, define $Irr^C(G) = \bigsqcup_{n \geq 0} Irr_n^C(G)$ (the disjoint topological sum).

We denote by $G^w = (G, w(G, Irr(G)))$ (resp. $G^{wC} = (G, w(G, Irr^C(G)))$) the group G equipped with the weak (group) topology generated by $Irr(G)$ (resp. $Irr^C(G)$). Since equivalent representations define the same topology, we have $G^w = (G, w(G, \widehat{G}))$. That is, the *weak topology* is the initial topology on G defined by the dual object. Moreover, in case G is a separable, metric, locally compact group, then every irreducible unitary representation acts on a separable Hilbert space and, as a consequence, is unitary equivalent to a member of $Irr^C(G)$. Thus $G^w = (G, w(G, Irr^C(G))) = G^{wC}$ for separable, metric, locally compact groups. We will make use of this fact in order to avoid the proliferation of isometries (see [14]). In case the group G is abelian, the dual object \widehat{G} is a group, which is called *dual group*, and the weak topology of G reduces

to the weak topology generated by all continuous homomorphisms of G into the unit circle \mathbb{T} . That is, the weak topology coincides with the so-called *Bohr topology* of G , that we recall next for the reader's sake.

With every (not necessarily abelian) topological group G there is associated a compact Hausdorff group bG , the so-called *Bohr compactification* of G , and a continuous homomorphism b of G onto a dense subgroup of bG such that bG is characterized by the following universal property: given any continuous homomorphism h of G into a compact group K , there is always a continuous homomorphism \bar{h} of bG into K such that $h = \bar{h} \circ b$ (see [28, V §4], where a detailed study on bG and their properties is given). In Anzai and Kakutani [3] bG is built when G is locally compact abelian (LCA). However, most authors agree that it was A. Weil [48] the first to build bG . Weil called bG “Groupe compact attaché à G ”. The name of Bohr compactification was given by Alfsen and Holm [2] in the context of arbitrary topological groups. The *Bohr topology* of a topological group G is the one that inherits as a subgroup of bG .

The weak topology of a topological group plays a role analogous to the weak topology in a Banach space. Therefore, it is often studied in connection to the original topology of the group. For instance, it can be said that the preservation of compact-like properties from G^w to G concerns “uniform boundedness” results and, in many cases, it can be applied to prove the continuity of certain related algebraic homomorphisms.

Our main result establishes that for every sequence $\{g_n\}_{n < \omega}$ in a locally compact group G , then either $\{g_n\}_{n < \omega}$ has a weak Cauchy subsequence or contains a subsequence that is an I_0 set. As a consequence, we obtain some sufficient conditions for the existence of (weak) Sidon sets in locally compact groups. It is still an open question

whether every infinite subset of a locally compact group G contains a (weak) Sidon subset (see [32, 20]).

As far as we know, the first result about the existence of I_0 sets was given by Hartman and Ryll-Nardzewski [25], who considered the weak topology associated to a locally compact abelian (LCA, for short) group and introduced the notion of *interpolation* (or I_0) set. As they defined it, a subset E of a LCA group G is an I_0 set if every bounded function on E is the restriction of an almost *periodic function* on G (here, it is said that a complex-valued function f defined on G is *almost periodic* when is the restriction of a continuous function defined on bG). Alternatively, one can define this notion without recurring to the Bohr compactification using the Fourier transform. Thus E is an I_0 set if every bounded function on E is the restriction of the Fourier transform of a discrete measure on G .

Therefore, an I_0 set is a subset E of G such that any bounded map on E can be interpolated by a continuous function on bG . As a consequence, if E is a countably infinite I_0 set, then \overline{E}^{bG} is canonically homeomorphic to $\beta\omega$. The main result given by Hartman and Ryll-Nardzewski is the following:

Theorem 1.1 ([25]). *Every LCA group G contains I_0 sets.*

For the particular case of discrete abelian groups, van Douwen achieved a remarkable progress by proving the existence of I_0 sets in very general situations. His main result can be formulated in the following way:

Theorem 1.2 ([45], Theorem 1.1.3). *Let G be a discrete Abelian group and let A be an infinite subset of G . Then there is a subset B of A with $|B| = |A|$ such that B is an I_0 set.*

In fact, van Douwen extended his result to the real line but left unresolved the question for LCA groups. In general, the weak topology of locally compact groups has been considered by many workers so far, specially for abelian groups, where the amount of important results in this direction is vast (see [24] for a recent a comprehensive source on the subject).

For those locally compact groups that can be injected in their Bohr compactification, the so-called *maximally almost periodic groups*, the existence of I_0 sets was clarified in [22]. However, many (non-abelian) locally compact groups cannot be injected in their Bohr compactification (that can become trivial in some cases). Thus, the question of extending Rosenthal's result to general locally compact groups remained open until now.

The starting point of our research lies on three celebrated results on $C(X)$, the space of continuous functions on a Polish space X , equipped with the pointwise convergence topology. The first two basic results are Rosenthal's dichotomy theorem [38] and a theorem by Bourgain, Fremlin and Talagrand about compact subsets of Baire class 1 functions [7], that we present in the way they are formulated by Todorčević in [42].

Theorem 1.3. (H. P. Rosenthal) *If X is a Polish space and $\{f_n\} \subseteq C(X)$ is a pointwise bounded sequence, then either $\{f_n\}$ contains a convergent subsequence or a subsequence whose closure in \mathbb{R}^X is homeomorphic to $\beta\omega$.*

Theorem 1.4. (J. Bourgain, D.H. Fremlin, M. Talagrand) *Let X be a Polish space and let $\{f_n\}_{n<\omega} \subseteq C(X)$ be a pointwise bounded sequence. The following assertions are equivalent (where the closure is taken in \mathbb{R}^X):*

(a) *$\{f_n\}_{n<\omega}$ is sequentially dense in its closure.*

(b) *The closure of $\{f_n\}_{n<\omega}$ contains no copy of $\beta\omega$.*

Our third starting fact is extracted from a result by Pol [37, p. 34], that again was formulated in different terms (cf. [8]). Here, we only use one of the implications established by Pol.

Theorem 1.5. (R. Pol) *Let X be a complete metric space, G a subset of $C(X)$ which is uniformly bounded. If $\overline{G}^{\mathbb{R}^X} \not\subseteq B_1(X)$, then G contains a sequence whose closure in \mathbb{R}^X is homeomorphic to $\beta\omega$.*

This paper is divided as follows: The first section is introductory in nature, recounting briefly the history of the topic and some basic results that are needed along the paper. In the second section, we study sets of continuous functions whose pointwise closure is compact and contained in the space of all Baire class 1 functions. We analyze the special case where the functions are defined from a polish space X to a metric space M . Rosenthal [38], Bourgain [6], and Bourgain, Fremlin and Talagrand [7] and, in a different direction, Todorćević [43] have extensively studied the compact subsets of $B_1(X)$. Our aim is to extend some of their fundamental results to the special case where the functions are metric-valued.

The result that connects with the rest of the paper is the extension of the Rosenthal's theorem [38]. Thus, the main goal is a dichotomy-type result for metric locally compact topological groups that has several applications in the study of general locally compact groups. For this purpose, we use the notion of I_0 set that plays a *rôle* analogous to the ℓ_1 -basis in the realm of locally compact groups.

In the fourth section, we look at Sidon subsets of non-abelian groups and we apply our previous results in order to obtain sufficient conditions for the existence of these interpolation sets. Here, a subset E of G is called (*weak*) *Sidon set* when every bounded function can be interpolated by a continuous function defined on the Eberlein compactification eG (defined below). This is a weaker property than the classical notion of *Sidon set* in general (see [36]) but both notions coincide for amenable groups.

To finish the paper we deal with the property of strongly respecting compactness introduced in [13]. Comfort, Trigos-Arrieta and Wu proved that every abelian locally compact group strongly respects compactness. Having in mind that Hughes [29] proved that every locally compact group, not necessarily abelian, respects compactness, we extend this result verifying that in fact every locally compact group strongly respects compactness. This improves previous results by Comfort, Trigos-Arrieta and Wu [13] and Galindo and Hernández [22] mentioned above.

We now formulate our main results. In the sequel wG will denote the weak compactification of a locally compact group G (defined below) and $inv(wG) \stackrel{\text{def}}{=} \{x \in wG : xy = yx = 1 \text{ for some } y \in wG\}$ will designate the group of units of wG .

Theorem A. Let (G, τ) be a metric locally compact group and let $\{g_n\}_{n < \omega}$ be a sequence in G . Then, either $\{g_n\}_{n < \omega}$ contains a weak Cauchy subsequence or an I_0 set.

Theorem B. Every non-precompact subset of a locally compact group whose closure is placed in $inv(wG)$ contains an infinite I_0 set.

Theorem C. Every non-precompact subset of a locally compact group whose closure is placed in $inv(wG)$ contains an infinite weak Sidon set.

Theorem D. Every locally compact group G strongly respects compactness.

2. BAIRE CLASS 1 FUNCTIONS

Let X be a topological space and let (M, d) be a metric space. We let M^X (resp. $C(X, M)$) denote the set of functions (resp. continuous functions) from X to M , equipped with the product (equivalently, pointwise convergence) topology, unless otherwise stated. We assume WLOG that $d(x, y) \leq 1$ for all x, y in M .

A function $f : X \rightarrow M$ is said to be *Baire class 1* if there is a sequence of continuous functions that converges pointwise to f . We denote by $B_1(X, M)$ the set of all M -valued Baire 1 functions on X . If $M = \mathbb{R}$ we simply write $B_1(X)$. A compact space K is called *Rosenthal compactum* if K can be embedded in $B_1(X)$ for some Polish space X . Let X be a topological space, the *tightness* of X , denoted $tg(X)$, is the smallest infinite cardinal κ such that for any subset $A \subseteq X$ and any point $x \in \overline{A}$ there is a subset $B \subseteq A$ with $|B| \leq \kappa$ and $x \in \overline{B}$. We write $[A]^{\leq \omega}$ to denote the set of all countable subsets of A .

Given a subset $G \subseteq C(X, M)$, it defines an equivalence relation on X by $x \sim y$ if and only if $g(x) = g(y)$ for all $g \in G$. If $\tilde{X} = X/\sim$ is the quotient space and $p : X \rightarrow \tilde{X}$ denotes the canonical quotient map, each $g \in G$ has associated a map $\tilde{g} \in C(\tilde{X}, M)$ defined as $\tilde{g}(\tilde{x}) \stackrel{\text{def}}{=} g(x)$ for any $x \in X$ with $p(x) = \tilde{x}$. Furthermore, if $\tilde{G} \stackrel{\text{def}}{=} \{\tilde{g} : g \in G\}$, we can extend this definition to the pointwise closure of \tilde{G} . Thus, each $g \in \overline{\tilde{G}}^{M^{\tilde{X}}}$ has associated a map $\tilde{g} \in \overline{\tilde{G}}^{M^{\tilde{X}}}$ such that $\tilde{g} \circ p = g$. We denote by X_G the topological space $(\tilde{X}, t_p(\tilde{G}))$. Note that X_G is metrizable if G is countable and it is Polish if X is compact and G is countable.

With the terminology introduced above, the following fact is easily verified (see [19]).

Fact 2.1. *If G be a subset of $C(X, M)$ such that \overline{G}^{M^X} is compact, then the map $p^* : M^{X_G} \rightarrow M^X$, defined by $p^*(\tilde{f}) = \tilde{f} \circ p$, is a homeomorphism of $\overline{G}^{M^{X_G}}$ onto \overline{G}^{M^X} .*

2.1. Real-valued Baire class 1 functions. A direct consequence of the strong results presented at the Introduction are the following corollaries whose simple proof is included for the reader's sake.

Corollary 2.2. *If X is a Polish space and G is a uniformly bounded subset of $C(X)$. Then $tg(\overline{G}^{\mathbb{R}^X}) \leq \omega$ if and only if $\overline{G}^{\mathbb{R}^X} \subseteq B_1(X)$.*

Proof. One implication is consequence of a well known result by Bourgain, Fremlin and Talagrand (cf. [42]). Therefore, assume that $tg(\overline{G}^{\mathbb{R}^X}) \leq \omega$. If $\overline{G}^{\mathbb{R}^X} \not\subseteq B_1(X)$, by Theorem 1.5, we can find a sequence $\{f_n\}_{n < \omega} \subseteq G$ whose closure in \mathbb{R}^X is canonically homeomorphic to $\beta\omega$. This implies that $\overline{G}^{\mathbb{R}^X}$ contains a copy of $\beta\omega$, which is a contradiction. \square

Corollary 2.3. *Let X be a Polish space, G a uniformly bounded subset of $C(X)$. The following assertions are equivalent:*

- (a) $tg(\overline{G}^{\mathbb{R}^X}) \leq \omega$.
- (b) $\overline{G}^{\mathbb{R}^X} \subseteq B_1(X)$.
- (c) G is sequentially dense in $\overline{G}^{\mathbb{R}^X}$.
- (d) $|\overline{G}^{\mathbb{R}^X}| \leq \mathfrak{c}$.
- (e) G does not contain any sequence whose closure in \mathbb{R}^X is homeomorphic to $\beta\omega$.

Proof. (a) \Leftrightarrow (b) is Corollary 2.2.

(b) \Rightarrow (c) was proved by Bourgain, Fremlin and Talagrand (see [42]).

$(b) \Rightarrow (d)$, $(a) \Rightarrow (e)$ and $(c) \Rightarrow (a)$ are obvious.

$(d) \Rightarrow (b)$ and $(e) \Rightarrow (b)$ are a consequence of Theorem 1.5.

□

Furthermore, according to results of Rosenthal [38] and Talagrand [41] we can also add the property of containing a sequence equivalent to the unit basis ℓ_1 .

Definition 2.4. Let $\{g_n\}_{n < \omega}$ be a uniformly bounded real (or complex) sequence of continuous functions on a set X . We say that $\{g_n\}_{n < \omega}$ is equivalent to the unit basis ℓ_1 if there exists a real constant $C > 0$ such that

$$\sum_{i=1}^N |a_i| \leq C \cdot \left\| \sum_{i=1}^N a_i g_i \right\|_{\infty}$$

for all scalars a_1, \dots, a_N and $N \in \omega$.

Theorem 2.5 (Talagrand [41]). *Let X be a compact and metric space and G a uniformly bounded subset of $C(X)$. The following assertions are equivalent:*

(a) $\overline{G}^{\mathbb{R}^X} \subseteq B_1(X)$.

(b) Every sequence in G has a weak-Cauchy subsequence.

(c) G does not contain any sequence equivalent to the ℓ_1 basis.

Remark 2.6. It is pertinent to notice here that using Rosenthal-Dor Theorem [15], Talagrand's result formulated above also holds for complex valued continuous functions.

A slight variation of Corollary 2.2 is also fulfilled if X is a compact space and G is countable.

Corollary 2.7. *Let X be a compact space and G a countable uniformly bounded subset of $C(X)$. Then $tg(\overline{G}^{\mathbb{R}^X}) \leq \omega$ if and only if G does not contain any sequence whose closure in \mathbb{R}^X is homeomorphic to $\beta\omega$.*

Proof. Let X_G be the quotient space associated to G equipped with the topology of pointwise convergence on G . According to Fact 2.1, we may assume WLOG that $X = X_G$ and therefore that is a Polish space. It now suffices to apply Corollary 2.3. \square

Corollary 2.8. *Let X be a compact space, G a countable uniformly bounded subset of $C(X)$. The following assertions are equivalent:*

- (a) $tg(\overline{G}^{\mathbb{R}^X}) \leq \omega$.
- (b) G does not contain any sequence whose closure in \mathbb{R}^X is homeomorphic to $\beta\omega$.
- (c) $\overline{G}^{\mathbb{R}^X}$ is a Rosenthal compactum.
- (d) $|\overline{G}^{\mathbb{R}^X}| \leq \mathfrak{c}$.
- (e) G does not contain any subsequence equivalent to the ℓ_1 basis.

Proof. If X_G denotes the quotient space associated to G , then Fact 2.1 implies that $\overline{G}^{\mathbb{R}^X}$ is canonically homeomorphic to $\overline{G}^{\mathbb{R}^{X_G}}$

(a) \Leftrightarrow (b) is Corollary 2.7.

(a) \Rightarrow (c) By Corollary 2.2, we have that $\overline{G}^{\mathbb{R}^{X_G}} \subseteq B_1(X_G)$. Thus $\overline{G}^{\mathbb{R}^{X_G}}$ and, consequently, also $\overline{G}^{\mathbb{R}^X}$ are Rosenthal compacta.

(c) \Rightarrow (d) and (d) \Rightarrow (b) are obvious.

(a) \Leftrightarrow (e) It follows from Fact 2.1, Corollary 2.2 and Theorem 2.5. \square

Corollary 2.9. *Let X be a compact space and G a uniformly bounded subset of $C(X)$.*

If $tg(\overline{G}^{\mathbb{R}^X}) \leq \omega$, then $\overline{G}^{\mathbb{R}^X} \subseteq B_1(X)$.

Proof. Suppose that there is $f \in \overline{G}^{\mathbb{R}^X} \setminus B_1(X)$. Since $tg(\overline{G}^{\mathbb{R}^X}) \leq \omega$, there is $L \in [G]^{\leq \omega}$ such that $f \in \overline{L}^{\mathbb{R}^X}$. Therefore $f \in \overline{L}^{\mathbb{R}^X} \setminus B_1(X)$ and, by Fact 2.1, we deduce that $\tilde{f} \in \overline{L}^{\mathbb{R}^{X_L}} \setminus B_1(X_L)$. It now suffices to apply Corollaries 2.3 and 2.7. \square

The following example shows that Corollary 2.8 may fail if one takes G of uncountable cardinality.

Example 2.10. Let $X = [0, \omega_1]$ and set

$$G = \{\chi_{[\alpha, \omega_1]} : \alpha < \omega_1, \text{ and } \alpha \text{ is not a limit ordinal}\}.$$

Then $|\overline{G}^{\mathbb{R}^X}| = \mathfrak{c}$. However $\chi_{\{\omega_1\}}$ is in the closure of G but it does not belong to the closure of any countable subset of G .

2.2. Metric-valued Baire class 1 functions. The goal in this section is to extend the results obtained for real-valued Baire class 1 functions to functions that take value in a metric space. This is accomplished using an idea of Christensen [12]. First, we need the following definition.

Let (M, d) be a metric space that we always assume equipped with a bounded metric.

We set

$$\mathcal{K} \stackrel{\text{def}}{=} \{\alpha : M \rightarrow [-1, 1] : |\alpha(m_1) - \alpha(m_2)| \leq d(m_1, m_2), \quad \forall m_1, m_2 \in M\}.$$

Being pointwise closed and equicontinuous by definition, it follows that \mathcal{K} is a compact and metrizable subspace of \mathbb{R}^M . For $m_0 \in M$, define $\alpha_{m_0} \in \mathbb{R}^M$ by $\alpha_{m_0}(m) \stackrel{\text{def}}{=} d(m, m_0)$ for all $m \in M$. It is easy to check that $\alpha_{m_0} \in \mathcal{K}$. Given $f \in M^X$ we associate a map $\check{f} \in \mathbb{R}^{X \times \mathcal{K}}$ defined by

$$\check{f}(x, \alpha) = \alpha(f(x)) \text{ for all } (x, \alpha) \in X \times \mathcal{K}.$$

In like manner, given any subset G of M^X we set $\check{G} \stackrel{\text{def}}{=} \{\check{f} : f \in G\}$.

Lemma 2.11. *Let X be a topological space, (M, d) a metric space and $G \subseteq C(X, M)$.*

Then:

- (a) $f \in C(X, M)$ if and only if $\check{f} \in C(X \times \mathcal{K})$.
- (b) A net $\{g_\delta\}_{\delta \in w} \subseteq C(X, M)$ converges to $f \in M^X$ if and only if the net $\{\check{g}_\delta\}_{\delta \in w} \subseteq C(X \times \mathcal{K})$ converges to $\check{f} \in \mathbb{R}^{X \times \mathcal{K}}$.
- (c) If \overline{G}^{M^X} is compact, then \overline{G}^{M^X} and $\overline{G}^{\mathbb{R}^{X \times \mathcal{K}}}$ are canonically homeomorphic.

Proof. (a) Suppose that $f \in C(X, M)$ and let $\{(x_\delta, \alpha_\delta)\}_{\delta \in w} \subseteq X \times \mathcal{K}$ be a net that converges to $(x, \alpha) \in X \times \mathcal{K}$. For every $\delta \in w$, we have

$$\begin{aligned} |\alpha_\delta(f(x_\delta)) - \alpha(f(x))| &\leq |\alpha_\delta(f(x_\delta)) - \alpha_\delta(f(x))| + |\alpha_\delta(f(x)) - \alpha(f(x))| \leq \\ &\leq d(f(x_\delta), f(x)) + |\alpha_\delta(f(x)) - \alpha(f(x))| \end{aligned}$$

Since $\{f(x_\delta)\}_{\delta \in w}$ converges to $f(x)$ and $\{\alpha_\delta\}_{\delta \in w}$ converges to α it follows that

$$\lim_{\delta \in w} \check{f}(x_\delta, \alpha_\delta) = \lim_{\delta \in w} \alpha_\delta(f(x_\delta)) = \alpha(f(x)) = \check{f}(x, \alpha).$$

Conversely, suppose that $\check{f} \in C(X \times \mathcal{K})$ and let $\{x_\delta\}_{\delta \in w} \subseteq X$ a net that converges to $x \in X$. Consider the map $\alpha_{f(x)} \in \mathcal{K}$. We have

$$\lim_{\delta \in w} d(f(x_\delta), f(x)) = \lim_{\delta \in w} \alpha_{f(x)}(f(x_\delta)) = \lim_{\delta \in w} \check{f}(x_\delta, \alpha_{f(x)}) = \check{f}(x, \alpha_{f(x)}) = 0,$$

That is, f is continuous.

(b) Suppose that $\{g_\delta\}_{\delta \in w} \subseteq C(X, M)$ converges pointwise to $f \in M^X$ and take the associated sequence $\{\check{g}_\delta\}_{\delta \in w} \subseteq C(X \times \mathcal{K})$. Then $\lim_{\delta \in w} \check{g}_\delta(x, \alpha) = \lim_{\delta \in w} \alpha(g_\delta(x)) = \alpha(\lim_{\delta \in w} g_\delta(x)) = \alpha(f(x)) = \check{f}(x, \alpha)$ for all $(x, \alpha) \in X \times \mathcal{K}$.

Conversely, suppose that $\{\check{g}_\delta\}_{\delta \in w} \subseteq C(X \times \mathcal{K})$ converges pointwise to $\check{f} \in \mathbb{R}^{X \times \mathcal{K}}$ and let us see that the sequence $\{g_\delta\}_{\delta \in w} \subseteq C(X, M)$ converges pointwise to f . Indeed,

it suffices to notice that for every $x \in X$ and its associated map $\alpha_{f(x)} \in \mathcal{K}$, we have

$$|\check{g}_\delta(x, \alpha_{f(x)}) - \check{f}(x, \alpha_{f(x)})| = |\alpha_{f(x)}(g_\delta(x)) - \alpha_{f(x)}(f(x))| = d(g_\delta(x), f(x)).$$

(c) Consider the map $\varphi : \overline{G}^{M^X} \rightarrow \overline{G}^{\mathbb{R}^{X \times \mathcal{K}}}$ defined by $\varphi(g) \stackrel{\text{def}}{=} \check{g}$ for all $g \in \overline{G}^{M^X}$. By compactness, it is enough to prove that φ is injective and continuous. The argument verifying the continuity of φ has been used in (b). Thus we only verify that φ is injective. Assume that $\varphi(f) = \varphi(g)$ with $f, g \in \overline{G}^{M^X}$, which means $\alpha(f(x)) = \alpha(g(x))$ for all $(x, \alpha) \in X \times \mathcal{K}$. Given $x \in X$, we have the map $\alpha_{g(x)} \in \mathcal{K}$ and, consequently, $\alpha_{g(x)}(f(x)) = \alpha_{g(x)}(g(x)) = 0$ for all $x \in X$. This yields $d(f(x), g(x)) = 0$ for all $x \in X$, which implies $f = g$. \square

Using the previous lemma we can generalize Theorem 1.3 to any metric space. This result will be very useful in the sequel.

Corollary 2.12. *Let X be a Polish space, (M, d) a metric space and $\{f_n\}_{n < \omega} \subseteq C(X, M)$ such that $\overline{\{f_n\}_{n < \omega}}^{M^X}$ is compact. Then, either $\{f_n\}_{n < \omega}$ contains a point-wise Cauchy subsequence or a subsequence whose closure in M^X is homeomorphic to $\beta\omega$.*

Now, we are in position of extending, to the setting of metric-valued functions, the results obtained in the previous section for real-valued functions.

Proposition 2.13. *Let X be a Polish space, (M, d) be a metric space and $G \subseteq C(X, M)$ such that \overline{G}^{M^X} is compact. The following assertions are equivalent:*

- (a) $tg(\overline{G}^{M^X}) \leq \omega$
- (b) $\overline{G}^{M^X} \subseteq B_1(X, M)$.
- (c) G is sequentially dense in \overline{G}^{M^X} .

(d) $|\overline{G}^{M^X}| \leq \mathfrak{c}$.

Proof. It follows from Lemma 2.11 and Corollary 2.3. □

Proposition 2.14. *Let X be a Polish space, (M, d) be a metric space and a sequence $\{f_n\}_{n < \omega} \subseteq C(X, M)$ such that $\overline{\{f_n\}_{n < \omega}}^{M^X}$ is compact. The following assertions are equivalent:*

- (a) $\{f_n\}_{n < \omega}$ is sequentially dense in its closure.
- (b) The closure of $\{f_n\}_{n < \omega}$ contains no copy of $\beta\omega$.

Proof. (a) \Rightarrow (b) is obvious.

(b) \Rightarrow (a) By Lemma 2.11 we know that $\overline{\{f_n\}_{n < \omega}}^{\mathbb{R}^X \times \mathbb{R}^X}$ contains no copy of $\beta\omega$. Thus by Theorem 1.4, it follows that $\{\check{f}_n\}_{n < \omega}$ is sequentially dense in its closure. By Lemma 2.11 we conclude that $\{f_n\}_{n < \omega}$ is sequentially dense in its closure. □

Corollary 2.15. *Let X be a compact space, (M, d) be a metric space $G \subseteq C(X, M)$ such that \overline{G}^{M^X} is compact. If $tg(\overline{G}^{M^X}) \leq \omega$, then $\overline{G}^{M^X} \subseteq B_1(X, M)$.*

Proof. It suffices to apply Corollary 2.9 and Lemma 2.11. □

Corollary 2.16. *Let X be a compact space, (M, d) be a metric space and G a countable subset of $C(X, M)$ such that \overline{G}^{M^X} is compact. The following assertions are equivalent:*

- (a) $tg(\overline{G}^{M^X}) \leq \omega$.
- (b) G does not contain any sequence whose closure in M^X is homeomorphic to $\beta\omega$.
- (c) \overline{G}^{M^X} is a Rosenthal compactum.
- (d) $|\overline{G}^{M^X}| \leq \mathfrak{c}$.

Proof. It suffices to apply Corollary 2.8 and Lemma 2.11. \square

Corollary 2.17. *Let X be a compact space, (M, d) be a metric space and G a countable subset of $C(X, M)$ such that \overline{G}^{M^X} is compact. If $|\overline{G}^{M^X}| \geq 2^{\mathfrak{c}}$, then there is a countable subset L of G such that its closure is canonically homeomorphic to $\beta\omega$.*

Proof. Use Corollary 2.16. \square

In the case where the metric space M is \mathbb{C} we have the following Corollary.

Corollary 2.18. *Let X be a compact space and G a uniformly and countable subset of $C(X, \mathbb{C})$. The following assertions are equivalent:*

- (a) $tg(\overline{G}^{\mathbb{C}^X}) \leq \omega$.
- (b) G does not contain any sequence whose closure in \mathbb{C}^X is homeomorphic to $\beta\omega$.
- (c) $\overline{G}^{\mathbb{C}^X}$ is a Rosenthal compactum.
- (d) $|\overline{G}^{\mathbb{C}^X}| \leq \mathfrak{c}$.
- (e) G does not contain a subsequence equivalent to the ℓ_1 basis.

Proof. Apply the complex version of Theorem 2.5 and Corollary 2.16. \square

3. DICHOTOMY-TYPE RESULT FOR LOCALLY COMPACT GROUPS

Recall from the Introduction that for a locally compact group G and if $\mathcal{H}_n \stackrel{\text{def}}{=} \mathbb{C}^n$ for $n = 1, 2, \dots$; $\mathcal{H}_0 \stackrel{\text{def}}{=} l^2(\mathbb{Z})$, then $\text{Irr}_n^C(G)$ denotes the set of irreducible unitary representations of G on \mathcal{H}_n (where it is assumed that every set $\text{Irr}_n^C(G)$ is equipped with the compact open topology), and $\text{Irr}^C(G) = \bigsqcup_{n \geq 0} \text{Irr}_n^C(G)$ (the disjoint topological sum).

The symbols G^w (resp. G^{wc}) designate the group G equipped with the weak (group) topology generated by $Irr(G)$ (resp. $Irr^C(G)$). As it was mentioned in the Introduction, if G is abelian, then the weak topology of G coincides with the so-called Bohr topology associated to G .

Definition 3.1. We denote by $P(G)$ the set of continuous positive definite functions on (G, τ) . If $\sigma \in Irr(G)$ and $v \in \mathcal{H}^\sigma$, then the positive definite function:

$$\varphi : g \mapsto \langle \sigma(g)(v), v \rangle, g \in G$$

is called *pure*, and the family of all such functions is denoted by $I(G)$. We also can define $I^C(G)$ the subset of $I(G)$ consisting of the elements whose irreducible representation is in $Irr^C(G)$. When G is abelian, the set $I(G)$ coincides with the dual group \widehat{G} of the group G .

The proof of the lemma below is straightforward.

Lemma 3.2. *Let G be a locally compact group. Then:*

- (a) $G^w = (G, w(G, I(G)))$.
- (b) $G^{wc} = (G, w(G, I^C(G)))$.

Remark 3.3. We recall that $G^w = G^{wc}$ if G is a separable, metrizable, locally compact group.

Definition 3.4. Let G be a locally compact group and consider the two following natural embeddings:

$$w : G \hookrightarrow \prod_{\varphi \in I(G)} \overline{\varphi(G)} \quad \text{and} \quad w_C : G \hookrightarrow \prod_{\varphi \in I^C(G)} \varphi(G)$$

$$w(g) = (\varphi(g))_{\varphi \in I(G)}$$

$$w_C(g) = (\varphi(g))_{\varphi \in I^C(G)}$$

We define the *weak compactification* wG (resp. *C-weak compactification* w_CG) of G as the pair (wG, w) (resp. (w_CG, w_C)), where $wG \stackrel{\text{def}}{=} \overline{w(G)}$ (resp. $w_CG \stackrel{\text{def}}{=} \overline{w_C(G)}$).

This compactification has been previously considered in [9, 10] using different techniques. Also Akemann and Walter [1] extended Pontryagin duality to non-abelian locally compact groups using the family of pure positive definite functions. Again, in case G is abelian, both compactifications, (wG, w) and (w_CG, w_C) , coincide with the Bohr compactification of G .

A better known compactification of a locally group G which is closely related to wG is defined as follows (cf. [17, 40]): let $\overline{B(G)}^{\|\cdot\|_\infty}$ denote the commutative C^* -algebra consisting of the uniform closure of the Fourier-Stieltjes algebra of G . Here, the Fourier-Stieltjes algebra is defined as the matrix coefficients of the unitary representations of G . Following [34] we call the spectrum eG of $\overline{B(G)}^{\|\cdot\|_\infty}$ the *Eberlein compactification* of G . Since the Eberlein compactification eG is defined using the family of all continuous positive definite functions, it follows that wG is a factor of eG and, as a consequence, inherits most of its properties. In particular, wG is a compact involutive semitopological semigroup.

The following definition was introduced by Hartman and Ryll-Nardzewski for abelian locally compact groups [25]. Here, we extend it to arbitrary not necessarily abelian locally compact groups.

Definition 3.5. A subset A of a locally compact group G is an I_0 set if every bounded complex (or real) valued function on A can be extended to a continuous function on wG . This definition extends the classic one, since when G is an abelian group, we have

that $wG = bG$, the so called *Bohr compactification* of G and $C(bG)|_G = AP(G)$ is the set of almost periodic functions on G .

Remark 3.6. Observe that if (G, τ) is a locally compact group and A be a countably infinite subset of G , then A is an I_0 set if and only if \overline{A}^{wG} is canonically homeomorphic to $\beta\omega$.

The following Lemma can be found in [42, Section 14, Th.3].

Lemma 3.7. *Let X be a compact space and $f : X \rightarrow \beta\omega$ a continuous and onto map. If $f^{-1}(n)$ is a singleton for all $n < \omega$ and $f^{-1}(\omega)$ is dense in X . Then f is a homeomorphism.*

Lemma 3.8. *Let (G, τ) be a separable metric locally compact group and $\{g_n\}_{n < \omega}$ be a sequence on G such that $\overline{\{g_n\}_{n < \omega}}^{w_C G} \cong \beta\omega$, then $\overline{\{g_n\}_{n < \omega}}^{wG} \cong \beta\omega$.*

Proof. Let $\varphi : G^w \rightarrow G^{w_C}$ be the identity map, which is clearly a continuous group homomorphism and set $\overline{\varphi} : wG \rightarrow w_C G$ the continuous extension of φ . The result follows from Lemma 3.7. \square

We now recall some known results about unitary representations of locally compact groups that are needed in the proof of our main result in this section. One main point is the decomposition of unitary representations by direct integrals of irreducible unitary representations. This was established by Mautner [33] following the ideas introduced by von Neuman in [47].

Theorem 3.9 (F. I. Mautner, [33]). *For any representation $(\sigma, \mathcal{H}_\sigma)$ of a separable locally compact group G , there is a measure space (R, \mathcal{R}, r) , a family $\{\sigma(p)\}$ of irreducible*

representations of G , which are associated to each $p \in R$, and an isometry U of \mathcal{H}_σ such that

$$U\sigma U^{-1} = \int_R \sigma(p) d_r(p).$$

Remark 3.10. The proof of the above theorem given by Mautner assumes that the representation space \mathcal{H}_σ is separable but, subsequently, Segal [39] removed this constraint. Furthermore, it is easily seen that we can assume that $\sigma(p)$ belongs to $\text{Irr}^C(G)$ locally almost everywhere in the theorem above (cf. [30]).

A remarkable consequence of Theorem 3.9 is the following corollary about positive definite functions.

Corollary 3.11. *Every Haar-measurable positive definite function φ on a separable locally compact group G can be expressed for all $g \in G$ outside a certain set of Haar-measure zero in the form*

$$\varphi(g) = \int_R \varphi_p(g) d_r(p),$$

where φ_p is a pure positive definite functions on G for all $p \in R$.

Definition 3.12. Let U be an open neighbourhood of the identity of a topological group G . We say that a sequence $\{g_n\}_{n < \omega}$ is U -discrete if $g_n U \cap g_m U = \emptyset$ for all $n \neq m \in \omega$.

Proof of Theorem A. Since G is metric, we may assume WLOG that the sequence is not $\{g_n\}_{n < \omega}$ is not τ -precompact. Otherwise, it would contain a τ -convergent subsequence that, as a consequence, would be weakly convergent and *a fortiori* weakly Cauchy.

Thus, $\{g_n\}_{n<\omega}$ must contain a subsequence that is U_0 -discrete for some symmetric, relatively compact and open neighbourhood of the identity U_0 in G . For simplicity's sake, we assume WLOG that the whole sequence $\{g_n\}_{n<\omega}$ is U_0 -discrete.

Take the σ -compact, open subgroup $H \stackrel{\text{def}}{=} \langle \overline{U}_0 \cup \{g_n\}_{n<\omega} \rangle$ of G . Since H is metric, σ -compact, it follows that H is a Polish locally compact group. Consequently, by [14, Section 18.1.10], we have that $\text{Irr}_m^C(H)$, equipped with the compact open topology, is a Polish space for all $m \in \{0, 1, 2, \dots\}$.

Being \mathcal{H}_m separable, for each $m \in \{0, 1, 2, \dots\}$, there exists a countable subset $D_m \stackrel{\text{def}}{=} \{v_n^m\}_{n<\omega}$ that is dense in the unit ball of \mathcal{H}_m (therefore, the linear subspace generated by D_m will be dense in \mathcal{H}_m). Fix $m \in \{0, 1, 2, \dots\}$ and let \mathbb{D} denote the closed unit disk in \mathbb{C} . We have that $\langle \sigma(g)(v_n^m), v_n^m \rangle \in \mathbb{D}$ for all $\sigma \in \text{Irr}_m^C(H)$, $g \in H$ and $n < \omega$.

For each $m \in \{0, 1, 2, \dots\}$, let $\alpha_m : H \rightarrow C_p(\text{Irr}_m^C(H), \mathbb{D}^\omega)$ be the continuous and injective map defined by $\alpha_m(h)(\sigma) \stackrel{\text{def}}{=} (\langle \sigma(h)(v_n^m), v_n^m \rangle)_{n \in \omega}$ for all $h \in H$ and $\sigma \in \text{Irr}_m^C(H)$. Since \mathbb{D}^ω is a compact metric space, it follows that

$$\overline{\{\alpha_m(g_n)\}_{n<\omega}}^{(\mathbb{D}^\omega)^{\text{Irr}_m^C(H)}}$$

is compact for all $m \in \{0, 1, 2, \dots\}$.

Now, we successively apply Corollary 2.12 for each $m \in \{0, 1, 2, \dots\}$ as follows.

For $m = 0$, $\{\alpha_0(g_n)\}_{n<\omega}$ contains either a pointwise Cauchy subsequence or a subsequence whose closure in $(\mathbb{D}^\omega)^{\text{Irr}_0^C(H)}$ is canonically homeomorphic to $\beta\omega$.

If there is a Cauchy subsequence $\{\alpha_0(g_{n_i^0})\}_{i<\omega}$, then we go on to the case $m = 1$. That is $\{\alpha_1(g_{n_i^0})\}_{i<\omega}$ contains either a pointwise Cauchy subsequence or a subsequence whose closure in $(\mathbb{D}^\omega)^{\text{Irr}_1^C(H)}$ is canonically homeomorphic to $\beta\omega$. If there is a Cauchy subsequence $\{\alpha_1(g_{n_i^1})\}_{i<\omega}$ we go on to the case $m = 2$, and so forth.

Assume that we can find a pointwise Cauchy subsequence in each step and take the diagonal subsequence $\{g_{n_i^i}\}_{i<\omega}$. We have that $\{\alpha_m(g_{n_i^i})\}_{i<\omega}$ is pointwise Cauchy for each $m \in \{0, 1, 2, \dots\}$. We claim that the subsequence $\{g_{n_i^i}\}_{i<\omega}$ is Cauchy in the weak topology of G .

Indeed, take an arbitrary element $\varphi \in I^C(H)$, then there is $t \in \{0, 1, 2, \dots\}$, $\sigma \in \text{Irr}_t^C(H)$ and $v \in \mathcal{H}_t$ such that $\varphi(h) = \langle \sigma(h)(v), v \rangle$ for all $h \in H$, where we may assume that $\|v\| \leq 1$ WLOG.

Let $\epsilon > 0$ be an arbitrary positive real number. By the density of D_t , there is $u \in D_t$ such that $\|u - v\| < \epsilon/6$.

For every $h \in H$, we have

$$\begin{aligned} | \langle \sigma(h)(v), v \rangle - \langle \sigma(h)(u), u \rangle | &= | \langle \sigma(h)(v), v \rangle - \langle \sigma(h)(u), v \rangle + \\ &\quad \langle \sigma(h)(u), v \rangle - \langle \sigma(h)(u), u \rangle | \\ &\leq | \langle \sigma(h)(v - u), v \rangle | + | \langle \sigma(h)(u), v - u \rangle | \\ &\leq 2\|v - u\| < \epsilon/3. \end{aligned}$$

On the other hand, we have that $\{\alpha_t(g_{n_i^i})\}_{i<\omega}$ is a pointwise Cauchy sequence in $(\mathbb{D}^\omega)^{\text{Irr}_t^C(H)}$. Thus, from the definition of α_t and, since $u \in D_t$, it follows that

$$\{ \langle \sigma(g_{n_i^i})(u), u \rangle \}_{i<\omega}$$

is a Cauchy sequence in \mathbb{D} . Hence, there is $i_0 < \omega$ such that

$$| \langle \sigma(g_{n_i^i})(u), u \rangle - \langle \sigma(g_{n_j^j})(u), u \rangle | < \epsilon/3 \text{ for all } i, j \geq i_0.$$

This yields

$$\begin{aligned}
| \langle \sigma(g_{n_i^i})(v), v \rangle - \langle \sigma(g_{n_j^j})(v), v \rangle | &\leq | \langle \sigma(g_{n_i^i})(v), v \rangle - \langle \sigma(g_{n_i^i})(u), u \rangle | \\
&+ | \langle \sigma(g_{n_i^i})(u), u \rangle - \langle \sigma(g_{n_j^j})(u), u \rangle | \\
&+ | \langle \sigma(g_{n_j^j})(u), u \rangle - \langle \sigma(g_{n_j^j})(v), v \rangle | \\
&< 3\epsilon/3 = \epsilon.
\end{aligned}$$

We conclude that $\{\langle \sigma(g_{n_i^i})(v), v \rangle\}_{i < \omega} = \{\varphi(g_{n_i^i})\}_{i < \omega}$ is a Cauchy sequence in \mathbb{D} for all $\varphi \in I^C(H)$. Since H is a locally compact Polish group, we have that $G^w = G^{wc}$ by Remark 3.3. As a consequence, it follows that which proves that $\{g_{n_i^i}\}_{i < \omega}$ is weakly Cauchy in H . We must now verify that $\{g_{n_i^i}\}_{i < \omega}$ is weakly Cauchy in G .

In order to do so, take a map $\psi \in I(G)$. Since H is separable, by Corollary 3.11, there is a measure space (R, \mathcal{R}, r) , a family $\{\psi_p\}$ of pure positive definite functions on H , which are associated to each $p \in R$, such that

$$\psi(h) = \int_R \psi_p(h) d_r(p) \text{ for all } h \in H.$$

Therefore

$$\psi(g_{n_i^i}) = \int_R \psi_p(g_{n_i^i}) d_r(p) \text{ for all } i < \omega.$$

Now, for each $i < \omega$, consider the map f_i on R by $f_i(p) \stackrel{\text{def}}{=} \psi_p(g_{n_i^i})$. Then f_i is integrable on R and, since $\{g_{n_i^i}\}_{i < \omega}$ is weakly Cauchy in H , it follows that $\{f_i\}$ is a pointwise Cauchy sequence on R . Furthermore, if $\psi_p(h) = \langle \sigma_p(h)[v_p], v_p \rangle$ for some $\sigma_p \in \text{Irr}(H)$ and $v_p \in \mathcal{H}_{\sigma_p}$, it follows that

$$|f_i(p)| = |\psi_p(g_{n_i^i})| = | \langle \sigma_p(g_{n_i^i})[v_p], v_p \rangle | \leq \|v_p\|^2.$$

Thus defining f on R as the pointwise limit of $\{f_i\}$, we are in position to apply Lebesgue's dominated convergence theorem in order to obtain that

$$\int_R f(p) d_r(p) = \lim_{i \rightarrow \infty} \int_R \psi_p(g_{n_i^i}) d_r(p) = \lim_{i \rightarrow \infty} \psi(g_{n_i^i}).$$

In other words, the sequence $\{\psi(g_{n_i^i})\}$ converges and, therefore, is Cauchy for all $\psi \in I(G)$. Hence $\{g_{n_i^i}\}$ is weakly Cauchy in G and we are done.

Suppose now that there exists an index $m_0 \in \{0, 1, 2, \dots, \infty\}$ such that $\{\alpha_{m_0}(g_{n_i^{m_0}})\}_{i < \omega}$ contains a subsequence $\{\alpha_{m_0}(g_{n(j)})\}_{j < \omega}$ whose closure in $(\mathbb{D}^\omega)^{Irr_{m_0}^C(H)}$ is homeomorphic to $\beta\omega$. Applying Lemma 3.7, we know that $\overline{\{g_{n(j)}\}_{j < \omega}}^{w_{CH}} \cong \beta\omega$. Consequently, by Lemma 3.8, we obtain that $\overline{\{g_{n(j)}\}_{j < \omega}}^{w_H} \cong \beta\omega$.

On the other hand, by Bichteler's [5, Lemma 3.2], we have that the irreducible representations of H are the restrictions of irreducible representations of G , which implies that the identity map $id : (H, w(G, I(G))|_H) \rightarrow (H, w(H, I(H)))$ is a continuous group isomorphism that can be extended canonically to a homeomorphism between their associated compactifications $\overline{id} : \overline{H}^{w_G} \rightarrow wH$. By Lemma 3.7 again, we obtain that $\overline{\{g_{n(j)}\}_{j < \omega}}^{w_G} \cong \beta\omega$. Thus $\{g_{n(j)}\}_{j < \omega}$ is an I_0 set, which completes the proof. \square

Remark 3.13. Theorem **A** fails if we try to extend it to every locally compact group or even to every compact group. Indeed, Fedorčuk [18] has proved that the existence of a compact space K of cardinality \mathfrak{c} without convergent sequences is compatible with **ZFC**. If we take the Bohr compactification of the free abelian group generated by K , then every sequence contained in K does not fulfil any of the two choices established in Rosenthal's dichotomy.

4. INTERPOLATION SETS

Hartman and Ryll-Nardzewski [25] proved that every abelian locally compact group contains an I_0 set. This result was improved in [21], where it was proved that every non-precompact subset of an abelian locally compact group contains an I_0 set. These sort of results do not hold for general locally compact groups unfortunately. Indeed, the Eberlein compactification of the group $SL_2(R)$ coincides with its one-point compactification, which means that each continuous positive definite function on $SL_2(R)$ converges at infinity (see [11]). Therefore, for this group, only the first case of the dichotomy result in Theorem **A** holds. If we search for interpolation sets, some extra conditions have to be assumed.

In this section we explore the application of the results in the previous sections in the study of interpolation sets in locally compact groups. First, we need the following result that was established by Ernest [16] (cf. [31]) for separable metric locally compact groups and convergent sequences and Subsequently extended for locally compact groups and compact subsets by Hughes [29] .

Proposition 4.1. (J. Ernest, J.R. Hughes) *Let (G, τ) be a locally compact group. Then (G, τ) and G^w contain the same compact subsets.*

In some special cases, Hughes' result implies the convergence of weakly Cauchy sequences.

Proposition 4.2. *Let (G, τ) be a locally compact group and suppose that $\{g_n\}_{n < \omega}$ is a Cauchy sequence in G^w . If $\overline{\{g_n\}_{n < \omega}}^{wG} \subseteq \text{inv}(wG)$, then $\{g_n\}_{n < \omega}$ is τ -convergent in G .*

Proof. Assume that $\{g_n\}_{n<\omega}$ is a Cauchy sequence in G^w . First, we verify that the sequence is a precompact subset of (G, τ) .

Indeed, we have that $\{g_n\}_{n<\omega}$ converges to some element $p \in \text{inv}(wG)$. If $\{g_n\}_{n<\omega}$ were not precompact in (G, τ) , there would be a neighbourhood of the neutral element U and a subsequence $\{g_{n(m)}\}_{m<\omega}$ such that $g_{n(m)}^{-1} \cdot g_{n(l)} \notin U$ for each $m, l < \omega$ with $m \neq l$. On the other hand, the sequence $\{g_{n(m)}^{-1} \cdot g_{n(m+1)}\}_{m<\omega}$ converges to $p^{-1}p$, the neutral element in G^w . This takes us to a contradiction because, by Proposition 4.1, it follows that $\{g_{n(m)}^{-1} \cdot g_{n(m+1)}\}_{m<\omega}$ must also converge to the neutral element in (G, τ) .

Therefore, the sequence $\{g_n\}_{n<\omega}$ is a precompact subset of (G, τ) . This implies that $p \in G$ and we are done. \square

Lemma 4.3. *Let (G, τ) be a locally compact group and let B be a non-precompact subset of G such that $\overline{B}^{wG} \subseteq \text{inv}(wG)$. Then there exist a open subgroup H of G , a compact and normal subgroup K of H , a quotient map $p : H \rightarrow H/K$ and a sequence $\{g_n\}_{n<\omega} \subseteq B \cap H$ such that H/K is a Polish group and $\overline{p(\{g_n\}_{n<\omega})}^{wH/K} \cong \beta\omega$.*

Proof. Since B is non-precompact there exists an open, symmetric and relatively compact neighbourhood of the identity U in G such that B contains a U -discrete sequence $\{g_n\}_{n<\omega}$.

Consider the subgroup $H \stackrel{\text{def}}{=} \langle \overline{U} \cup \{g_n\}_{n<\omega} \rangle$, which is σ -compact and open in G . By Kakutani-Kodaira's theorem, there exists a normal, compact K of H such that $K \subseteq U$ and H/K is metrizable, and consequently Polish. Let $p : H \rightarrow H/K$ be the quotient map and let $\overline{p} : wH \rightarrow wH/K$ denote the canonical extension to the weak compactifications. Therefore, we have that $\overline{p}(\text{inv}(wH)) \subseteq \text{inv}(wH/K)$. Furthermore, since \overline{H}^{wG}

is canonically homeomorphic to wH , it follows that $\overline{\{g_n\}_{n<\omega}}^{wH} \subseteq \text{inv}(wH)$. Hence $\overline{\{p(g_n)\}_{n<\omega}}^{wH/K} \subseteq \text{inv}(wH/K)$. Thus, we are in position of applying Proposition 4.2.

Assume that there is a weakly Cauchy subsequence $\{p(g_s)\}_{s<\omega}$ in $(H/K)^w$, which would be τ/K -convergent by Proposition 4.2. Then by a theorem of Varopoulos [46], the sequence $\{p(g_s)\}_{s<\omega}$ could be lifted to a sequence $\{x_s\}_{s<\omega} \subseteq H$ converging to some point $x_0 \in H$. This would entail that $x_s^{-1}g_s \in K$ for all $s \in \omega$. Thus the sequence $\{g_s\}_{s<\omega}$ would be contained in the compact subset $(\{x_s\}_{s<\omega} \cup \{x_0\})K$, which is a contradiction since $\{g_n\}_{n<\omega}$ was supposed to be U -discrete. This contradiction completes the proof. \square

Proof of Theorem B. Apply Lemmata 3.7 and 4.3. \square

Corollary 4.4. *Let G be a discrete group and let $\{g_n\}_{n<\omega}$ be an infinite sequence in G . If $\overline{\{(g_n)\}_{n<\omega}}^{wG} \subseteq \text{inv}(wG)$, then $\{g_n\}_{n<\omega}$ contains an infinite I_0 set.*

Remark 4.5. In case the group G is abelian, Corollary 4.4 is a variant of van Douwen's Theorem 1.2.

We now look at *Sidon sets*, a well known family of interpolation sets in harmonic analysis. We notice that the following question still remains open (see [32] and [20, p. 57]).

Question 4.6 (Figà-Talamanca, 1977). *Do infinite weak Sidon exist in every infinite discrete non-abelian group?*

Since every I_0 set is automatically Sidon, Theorem B and Corollary 4.4 gives sufficient conditions for the existence of weak Sidon sets.

Definition 4.7. Let G be a group. A sequence $\{x_n\}_{n<\omega} \subseteq G$ is called *independent* if for every $n_0 < \omega$ the element $x_{n_0} \notin \langle \{x_n\}_{n \in \omega \setminus n_0} \rangle$. A group G is called *locally finite* if every finite subset of the group generates a finite subgroup. The group G is *residually finite* if for every non-identity element g of G there exists a normal subgroup N of finite index in G such that $g \notin N$. Finally, the group G is called an *FC-group* if every conjugacy class of G is finite.

Proposition 4.8. *Every independent sequence in a discrete group G is a weak Sidon set.*

Proof. Let $E = \{x_n\}_{n<\omega}$ be an independent sequence in G . It will suffice to show that \overline{E}^{eG} is homeomorphic to $\beta\omega$ or, equivalently, that every pair of disjoint subsets in E have disjoint closures in eG .

Indeed, for any pair A, B of arbitrary disjoint subsets of ω , set $X_A \stackrel{\text{def}}{=} \{x_n\}_{n \in A}$, and $X_B \stackrel{\text{def}}{=} \{x_n\}_{n \in B} \subseteq G \setminus \langle X_A \rangle$. Since $\langle X_A \rangle$ is an open subgroup of G , the positive definite function h defined by $h(x) = 1$ if $x \in \langle X_A \rangle$ and $h(x) = 0$ if $x \in G \setminus \langle X_A \rangle$ is continuous (see [27, 32.43(a)]). Thus $\overline{X_A}^{eG} \cap \overline{X_B}^{eG} = \emptyset$, which completes the proof. \square

Corollary 4.9. *Let G be a discrete group and let $\{x_n\}_{n<\omega}$ be a sequence in G . If the sequence contains either an independent subsequence or its wG -closure is contained in $\text{inv}(wG)$, then the sequence contains a weak Sidon set.*

Remark 4.10. Note that if G is a discrete *FC-group* we can find an independent sequence within every infinite subset of G . Therefore, every infinite subset of a discrete *FC-group* contains a weak Sidon set [35].

We finish this section with an example of a sequence that is a weak Sidon set and converges to the neutral element in the Bohr topology.

Proposition 4.11. *Let G be a residually finite, locally finite group that is not abelian by finite. Then G contains a sequence that is weak Sidon and converges in the Bohr topology.*

Proof. Let a_1 be a non trivial element in G' . Then, there is a finite non Abelian subgroup B_1 of G with $a_1 \in B'_1$. Suppose we have defined $L = \sum_{i=1}^n B_i$. Since L is finite and G is residually finite, we can get $m \in \mathbb{N}$ and $\sigma \in \text{Rep}_m(G)$ such that $\sigma|_L$ is faithful and $\sigma(G)$ is finite. Set $N = \ker \sigma$. If N were Abelian, it would follow that G is Abelian by finite, which is impossible. Thus, we may assume WLOG that $N' \neq \{1\}$ and we can replace G by N in order to obtain a finite non Abelian subgroup B_{n+1} of N and a non trivial element $a_{n+1} \in B'_{n+1}$. Using an inductive argument, we obtain $B = \sum_{i=1}^{\infty} B_i$, a subgroup of G , such that $B'_i \neq \{1\}$, for all $i \in \mathbb{N}$. Under such circumstances, it is known that the sequence $\{a_n\}$ is convergent in the Bohr topology (cf. [26, Cor. 3.10]). On the other hand, the sequence $\{a_n\}$ is an independent set by definition and, therefore, a weak Sidon subset in G . This completes the proof. \square

Remark that a positive answer to the following question will also solve Question 4.6.

Problem 4.12. *Does every discrete group contain an infinite I_0 set?*

5. PROPERTY OF STRONGLY RESPECTING COMPACTNESS

One of the first papers dealing with the weak topology of locally compact groups was given by Glicksberg in [23], where he proved that every weakly compact subset of a locally compact abelian group must be compact. Using the terminology introduced by Trigos-Arrieta in [44], it is said that locally compact abelian groups *respect compactness*. Along this line, Comfort, Trigos-Arrieta and Wu [13] established the following substantial extension of Glicksberg's theorem: Let G be a locally compact

abelian group with Bohr compactification bG and let N be a metrizable closed subgroup of bG . Then a subset A of G satisfies that $AN \cap G$ is compact in G if and only if AN is compact in bG . Motivated by this result, the authors defined a group that *strongly respects compactness* to be a group that satisfies the assertion above. They also raised the question of characterizing such groups. This question was considered in [22] but looking at the finite dimensional, irreducible representations of a locally compact group. Here we consider it in complete generality. First, we need the following definition that extends the notion of strongly respecting compactness to groups that are not necessarily abelian equipped with the weak topology.

Definition 5.1. We say that a locally compact group G *strongly respects compactness* if for any closed metrizable subgroup N of $\text{inv}(\text{w}G)$, a subset A of G satisfies that $AN \cap G$ is compact in G if and only if AN is compact in $\text{w}G$.

The main result of this section states that every locally compact group strongly respects compactness.

Let G be a locally compact group and let H be an open subgroup of G . We have discussed in the proof of Theorem A that the identity map $\text{id} : (H, \text{w}(G, I(G))|_H) \rightarrow H^{\text{w}}$ is a continuous isomorphism that can be extended to a continuous map $\overline{\text{id}} : \overline{H}^{\text{w}G} \rightarrow \text{w}H$ defined on their respective compactifications. On the other hand, using that $\text{w}G$ is a factor of the Eberlein compactification eG , it follows that $\text{w}G$ is a compact involutive semitopological semigroup for every locally compact group G . Taking this fact into account, the following lemma is easily verified.

Lemma 5.2. *Let G and H be locally compact groups and let $h : G \rightarrow H$ be a continuous homomorphism. Then there is a canonical continuous extension $\overline{h} : \overline{H}^{\text{w}G} \rightarrow \text{w}H$ such that for every p, q in $\overline{H}^{\text{w}G}$, we have $\overline{h}(pq) = \overline{h}(p)\overline{h}(q)$.*

The next result is a variation of [22, Lemma 3.6].

Lemma 5.3. *Let G be a locally compact group, H an open subgroup of G , A a subset of G , and let N be a subgroup of $\text{inv}(\text{w}G)$, containing the identity, such that AN is compact in $\text{w}G$. If F is an arbitrary subset of $AN \cap H$, then there exists $A_0 \subseteq A$ with $|A_0| \leq |N|$ such that*

$$\overline{F}^{\text{w}H} \subseteq \overline{(FF^{-1})}^{H^{\text{w}}} \cdot \overline{\text{id}}(A_0N).$$

Proof. We first verify that $\overline{F}^{\text{w}H} \subseteq \overline{\text{id}}(AN \cap \overline{H}^{\text{w}G})$.

Indeed, since $AN \cap H \subseteq AN \cap \overline{H}^{\text{w}G}$ and $AN \cap \overline{H}^{\text{w}G}$ is compact, we have $AN \cap H \subseteq \overline{\text{id}}(AN \cap \overline{H}^{\text{w}G})$ and, as a consequence, it follows that $\overline{AN \cap H}^{\text{w}H} \subseteq \overline{\text{id}}(AN \cap \overline{H}^{\text{w}G})$. Hence $\overline{F}^{\text{w}H} \subseteq \overline{\text{id}}(AN \cap \overline{H}^{\text{w}G})$ and $\overline{F}^{\text{w}H}$ is compact.

For any $x \in N$ such that $\overline{F}^{\text{w}H} \cap \overline{\text{id}}(Ax) \neq \emptyset$, pick $a_x \in A$ with $\overline{\text{id}}(a_x x) \in \overline{F}^{\text{w}H}$. We define $A_0 \stackrel{\text{def}}{=} \{a_x \in A : x \in N \text{ and } \overline{\text{id}}(a_x x) \in \overline{F}^{\text{w}H}\}$. We have $A_0 \subseteq A$ and $|A_0| \leq |N|$.

Pick an arbitrary point $b \in \overline{F}^{\text{w}H}$. Since $\overline{F}^{\text{w}H} \subseteq \overline{\text{id}}(AN \cap \overline{H}^{\text{w}G})$ we can find $a \in A$ and $y \in N$ such that $b = \overline{\text{id}}(ay)$. Set $b' = \overline{\text{id}}(a_y y) \in \overline{F}^{\text{w}H}$. Then $bb'^{-1} = \overline{\text{id}}(ay)\overline{\text{id}}(a_y y)^{-1} \in \overline{F}^{\text{w}H}\overline{F}^{-1\text{w}H} = \overline{FF^{-1}}^{\text{w}H}$. Observe also that, by Lemma 5.2, we have $bb'^{-1} = \overline{\text{id}}(ayy^{-1}a_y^{-1}) = \overline{\text{id}}(aa_y^{-1}) = aa_y^{-1} \in \text{w}H \cap G = H$. Therefore $bb'^{-1} \in \overline{FF^{-1}}^{\text{w}H} \cap H$. Since H is an open subgroup, by [29, Cor. 14.2], we deduce that $bb'^{-1} \in \overline{FF^{-1}}^{\text{w}H} \cap H = \overline{FF^{-1}}^{H^{\text{w}}}$. Thus $b = bb'^{-1}b' \in \overline{FF^{-1}}^{H^{\text{w}}} \cdot \overline{\text{id}}(A_0N)$ and we are done. \square

Proof of Theorem D. Let N be a metrizable subgroup of $\text{inv}(\text{w}G)$ and let A a subset of G such that AN is compact in $\text{w}G$. Since $AN \cap G$ is closed in G it suffices to see that it is precompact. By reduction to absurd, assume that $AN \cap G$ is non-precompact. By Theorem 4.3 there exist an open subgroup H of G , a compact and

normal subgroup K of H , a quotient map $p : H \rightarrow H/K$ and a sequence $F \subseteq AN \cap H$ such that H/K is a Polish group and $\overline{p(F)}^{wH/K} \cong \beta\omega$. Thus, $|\overline{p(F)}^{wH/K}| \geq 2^{\mathfrak{c}}$.

By Lemma 5.3 there is $A_0 \subseteq A$ with $|A_0| \leq |N|$ such that $\overline{F}^{wH} \subseteq \overline{(FF^{-1})}^{H^w} \overline{id}(A_0N)$. Since $|\overline{id}(A_0N)| \leq \mathfrak{c}$ we can enumerate it as $\{a_\alpha\}_{\alpha < \mathfrak{c}}$. Therefore, we can write $\overline{F}^{wH} \subseteq \bigcup_{\alpha < \mathfrak{c}} \overline{FF^{-1}}^{H^w} a_\alpha$.

Let $\overline{p} : wH \rightarrow wH/K$ be the canonical extension of p to the respective compactifications of H and H/K . Using Lemma 5.2, for each $z \in wH$ consider the map T_z defined on wH/K by $T_z(\overline{p}(x)) = \overline{p}(xz) = \overline{p}(x)\overline{p}(z)$ for all $x \in wH$. Hence, from the previous inclusion we obtain that $\overline{p(F)}^{wH/K} = \overline{p}(\overline{F}^{wH}) \subseteq \bigcup_{\alpha < \mathfrak{c}} T_{a_\alpha}(\overline{p(FF^{-1})}^{H^w/K})$. Since the topology of H^w/K is finer than that of $(H/K)^w$ we have that $\overline{p(FF^{-1})}^{H^w/K} \subseteq \overline{p(FF^{-1})}^{(H/K)^w}$. Furthermore $|\overline{p(FF^{-1})}^{(H/K)^w}| \leq \mathfrak{c}$ because H/K is a Polish space. Therefore, $|\overline{p(F)}^{wH/K}| \leq \mathfrak{c}$. This is a contradiction that completes the proof. \square

REFERENCES

- [1] Charles A. Akemann and Martin E. Walter. Non-abelian Pontriagin duality. *Duke Math. J.*, 39:451–463, 1972.
- [2] EM Alfsen and P Holm. A note on compact representations and almost periodicity in topological groups. *Mathematica Scandinavica*, 1962.
- [3] H Anzai and S Kakutani. Bohr Compactifications of a Locally Compact Abelian Group I. *Proceedings of the Imperial Academy*, 1943.
- [4] Bachir Bekka, Pierre de la Harpe, and Alain Valette. *Kazhdan's property*. Cambridge: Cambridge University Press, 2008.
- [5] K Bichteler. Locally compact topologies on a group and the corresponding continuous irreducible representations. *Pacific Journal of Mathematics*, 31:583–593, 1969.
- [6] J Bourgain. Compact sets of first baire class. *Bull. Soc. Math. Belg.*, 29(2):135–143, 1977.
- [7] J Bourgain, DH Fremlin, and M Talagrand. Pointwise compact sets of Baire-measurable functions. *American Journal of Mathematics*, 100(4):845–886, 1978.
- [8] B Cascales, I Namioka, and G Vera. The Lindelöf property and fragmentability. *Proc. Amer. Math. Soc.*, 128(11):3301–3309, 2000.
- [9] Michael Yin-Hei Cheng. Subalgebras generated by extreme points in Fourier-Stieltjes algebras of locally compact groups. *Stud. Math.*, 202(3):289–302, 2011.
- [10] Yin-Hei Cheng, Brian E. Forrest, and Nico Spronk. On the subalgebra of a Fourier-Stieltjes algebra generated by pure positive definite functions. *Monatsh. Math.*, 171(3-4):305–314, 2013.
- [11] Ching Chou. Minimally weakly almost periodic groups. *J. Funct. Anal.*, 36:1–17, 1980.

- [12] JPR Christensen. Joint Continuity of separately continuous functions. *Proceedings of the American Mathematical Society*, 82(3):455–461, 1981.
- [13] W.W. Comfort, F.J. Trigos-Arrieta, and T.-S. Wu. The Bohr compactification , modulo a metrizable subgroup. *Fund. Math.*, 143:119–136, 1993.
- [14] J Dixmier. C^* -Algèbres et leurs représentations. *Gauthier-Villars & Cie, Éditeur-Imprimeur, Paris*, 1964.
- [15] Leonard E. Dor and Haskell P. Rosenthal. On sequences spanning a complex ℓ_1 space. *Proceedings of the American Mathematical Society*, 47(2):515–515, feb 1975.
- [16] J Ernest. A strong duality theorem for separable locally compact groups. *Transactions of the American Mathematical Society*, 156:287–307, 1971.
- [17] P Eymard. L’algèbre de Fourier d’un groupe localement compact. *Bulletin de la Société Mathématique de France*, 92:181–236, 1964.
- [18] VV Fedorčuk. A compact space having the cardinality of the continuum with no convergent sequences. *Math. Proc. Cambridge Philos. Soc.*, 81(2):177–181, 1977.
- [19] M. Ferrer, S. Hernández, and L. Tárrega. Equicontinuity criteria for metric-valued sets of continuous functions. *Topology and its Applications*, Pending.
- [20] Alessandro Figa-Talamanca. Insiemi lacunari nei gruppi non commutativi. *Rend. Semin. Mat. Fis. Milano*, 47:45–59, 1977.
- [21] J Galindo and S Hernández. The concept of boundedness and the Bohr compactification of a MAP abelian group. *Fundamenta Mathematicae*, 159(3):195–218, 1999.
- [22] J Galindo and S Hernández. Interpolation sets and the Bohr topology of locally compact groups. *Advances in Mathematics*, 188:51–68, 2004.
- [23] I Glicksberg. Uniform boundedness for groups. *Canad. J. Math.*, 14:269–276, 1962.
- [24] C Graham and KE Hare. Interpolation and Sidon sets for compact groups. 2013.
- [25] S Hartman and C Ryll-Nardzewski. Almost periodic extensions of functions. *Colloquium Mathematicae*, 12(1):29–39, 1964.
- [26] S. Hernández and Ta-Sun Wu. Some new results on the Chu duality of discrete groups. *Monatsh. Math.*, 149(3):215–232, 2006.
- [27] Edwin Hewitt and Kenneth A. Ross. *Abstract harmonic analysis. Volume II, Structure and Analysis for Compact Groups Analysis on Locally Compact Abelian Groups*. Springer, 1970.
- [28] H. Heyer. *Dualität lokalkompakter Gruppen*. Berlin-Heidelberg-New York: Springer-Verlag, 1970.
- [29] JR Hughes. Weak compactness and topological groups. *Thesis (Ph.D.)The University of North Carolina at Chapel Hill*, 1972.
- [30] K Ikeshoji and Y Nakagami. On a strong duality for separable locally compact groups. *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 33(2):377–389, 1979.
- [31] K Ikeshoji and Y Nakagami. On a strong duality for separable locally compact groups. *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 33(2):377–389, 1979.
- [32] Jorge M. Lopez and Kenneth A. Ross. Sidon sets. *Lecture Notes in Pure and Applied Mathematics*. Vol. 13. New York: Marcel Dekker, Inc. V, 193 p. SFr. 50.00 (1975)., 1975.
- [33] F.I. Mautner. Unitary representations of locally compact groups. II. *Ann. Math. (2)*, 52:528–556, 1950.
- [34] M. Mayer. Strongly mixing groups. *Semigroup Forum*, 54(3):303–316, 1997.
- [35] Leonede De Michele and Paolo M. Soardi. Existence of Sidon Sets in Discrete FC-Groups. *Proceedings of the American Mathematical Society*, 55(2):457, mar 1976.
- [36] Massimo Angelo Picardello. Lacunary sets in discrete noncommutative groups. *Boll. Unione Mat. Ital., IV. Ser.*, 8:494–508, 1973.
- [37] R Pol. On pointwise and weak topology in function spaces. 1984.

- [38] HP Rosenthal. A characterization of Banach spaces containing ℓ_1 . *Proc. Nat. Acad. Sci. USA*, 71(6):2411–2413, 1974.
- [39] I.E. Segal. Decompositions of operator algebras I and II. *Mem. Am. Math. Soc.*, 9:67, 1951.
- [40] Nico Spronk and Ross Stokke. Matrix coefficients of unitary representations and associated compactifications. *Indiana Univ. Math. J.*, 62(1):99–148, 2013.
- [41] M Talagrand. Pettis integral and measure theory. 1984.
- [42] S Todorćević. Topics in topology. 1997.
- [43] Stevo Todorćević. Compact subsets of the first Baire class. *J. Am. Math. Soc.*, 12(4):1179–1212, 1999.
- [44] F.J. Trigos-Arrieta. Pseudocompactness on groups. *Ph.D Thesis, Wesleyan University*, 1991.
- [45] EK van Douwen. The maximal totally bounded group topology on G and the biggest minimal G -space, for abelian groups G . *Topology and its Applications*, 34(1):64–91, 1990.
- [46] NT Varopoulos. A theorem on the continuity of homomorphisms of locally compact groups. *Proc. Cambridge Philos. Soc.*, 60:449–463, 1964.
- [47] John von Neumann. On rings of operators. Reduction theory. *Ann. Math. (2)*, 50:401–485, 1949.
- [48] A Weil. L'intégration dans les groupes topologiques et ses applications. 1965.

UNIVERSITAT JAUME I, INSTITUTO DE MATEMÀTICAS DE CASTELLÓN, CAMPUS DE RIU SEC,
12071 CASTELLÓN, SPAIN.

E-mail address: mferrer@mat.uji.es

UNIVERSITAT JAUME I, DEPARTAMENTO DE MATEMÀTICAS, CAMPUS DE RIU SEC, 12071 CASTELLÓN,
SPAIN.

E-mail address: hernande@mat.uji.es

UNIVERSITAT JAUME I, IMAC AND DEPARTAMENTO DE MATEMÀTICAS, CAMPUS DE RIU SEC,
12071 CASTELLÓN, SPAIN.

E-mail address: ltarrega@uji.es